

CURRENT ALGEBRAS IN 3 + 1 DIMENSIONS

JOUKO MICKELSSON

Department of Mathematics, University of
Jyväskylä, SF-40100, Jyväskylä, Finland

ABSTRACT Aspects of a generalized representation theory of current algebras in 3+1 dimensions are discussed in terms of the Fock bundle method, the sesquilinear form approach (of Langmann and Ruijsenaars), and Hilbert space cocycles.

Let G be a Lie group of symmetries of a classical relativistic field theory. If $\{T_1, \dots, T_N\}$ is a basis for the Lie algebra $\mathfrak{g}$ of G then to each T_a there corresponds, by Noether's theorem, a conserved current $j_\mu^a(x)$. (Actually, the current should be thought of as a 3-form.)

When the classical field theory is quantized (in the Hamiltonian approach) on a space-like surface S then to each current j^a is supposed to correspond an operator valued distribution on the surface S . Formally,

$$(1) \quad j^a(f) = \int_S f(x) j^a(x),$$

where f is a smooth test function. Naively, one would expect that the charges $j^a(f)$ satisfy the commutation relations

$$(2) \quad [j^a(f), j^b(g)] = \lambda_c^{ab} j^c(fg),$$

where the λ_c^{ab} 's are the structure constants, $[T_a, T_b] = \lambda_c^{ab} T_c$.

However, in order to make sense of the quantum operators $j^a(f)$ one normally has to do certain regularizations. For example, in a field theory involving fermions in 1+1 space-time dimensions, transforming according to a finite-dimensional representation ρ of G , the group generated by the charge densities is the group $Map(S, G)$ of smooth maps from S to G , acting on the fermion field ψ by pointwise multiplications $\psi'(x) = \rho(f(x))\psi(x)$. In the second quantized theory the charge densities are given naively by the operators

$$(3) \quad T(f) = \sum f_{nm} a_n^* a_m,$$

where the a_n^* 's are the fermion creation operators and the a_m 's are annihilation operators; here f_{nm} denote the matrix elements of the multiplication operator in a given orthonormal basis in the one-particle space.

The problem with the the above construction is that in general a state created from the vacuum would have infinite norm. For making the operators finite it is sufficient to make a normal ordering: The products $a_n^* a_m$ are replaced by $: a_n^* a_m :$ where the dots mean that the operators lowering the energy are placed to the right. The normal ordering changes the commutation relations (2). Let π_+ be the projection onto the positive energy subspace in the one-particle Hilbert space H and $\epsilon = 2\pi_+ - 1$. Denoting by X, Y components of the charges in the one-particle representation, the new commutators are

$$(4) \quad [X, Y]' = [X, Y] + \frac{1}{4} \text{tr} \epsilon [[\epsilon, X], [\epsilon, Y]].$$

The second term on the right in (4) is an antisymmetric bilinear form c on a Lie algebra $\mathfrak{gl}_1$ which contains the algebra of integrated charge densities $j^a(f)$. The algebra $\mathfrak{gl}_1$ consists of bounded operators X in H such that $[\epsilon, X]$ is a Hilbert-Schmidt operator. The form c (Lundberg's cocycle, [Lu]) satisfies the *cocycle condition*

$$(5) \quad c(X, [Y, Z]) + c(Y, [Z, X]) + c(Z, [X, Y]) = 0$$

which guarantees that the Jacobi identity for the modified commutators $[\cdot, \cdot]'$ is satisfied.

In quantum field theory in space-time dimensions higher than two the normal ordering of the charge densities is not sufficient to make them finite; more radical regularizations are needed. In fact, there is no meaningful way to make the $j^a(f)$'s into operators in the (fermionic or bosonic) Fock space. The technical reason for this is that the currents are no more contained in $\mathfrak{gl}_1$ but in a bigger Lie algebra $\mathfrak{gl}_p$, where p must be greater or equal to $\frac{1}{2}(d+1)$, where d is the space dimension and $\mathfrak{gl}_p$ is characterized by the property that $[\epsilon, X]$ belongs to the Schatten ideal L_{2p} consisting of operators T such that $(T^*T)^p$ has a finite trace. Note that the cocycle c is defined only on $\mathfrak{gl}_1$ and not on any of the larger algebras $\mathfrak{gl}_p$, $p > 1$.

There is a modified cocycle c_p which converges for $\mathfrak{gl}_p$. The coefficient ring for these cocycles is not $\mathbb{C}$ but a certain space of complex valued functions on an infinite-dimensional manifold (which in this talk will be a Grassmannian manifold). For example, when $d = 3$ the cocycle takes the form, [MR],

$$(6) \quad c_2(X, Y; F) = \frac{1}{8} \text{tr}(\epsilon - F)[[\epsilon, X], [\epsilon, Y]],$$

where F is a hermitean operator, with $F^2 = 1$ and $\epsilon F + F \epsilon - 2\epsilon \in L_2$, parametrizing points on a Grassmannian Gr_2 . For $p > 2$ the cocycles c_p have been computed in [FT].

The new Lie algebra $\widehat{\mathfrak{gl}}_2$ defined by the cocycle (6) does not have interesting Hilbert space representations, [P]. Thus we have to look for a generalization of representation theory.

Because of the "background field" F the current algebra acts in a bundle of Fock spaces parametrized by Gr_2 , [M1]. There is an alternative description of the generalized representation of the current algebra proposed by E. Langmann [L]. The starting point is the observation of S.N.M. Ruijsenaars that although the charges are not well-defined operators in the Fock space, they still make sense as sesquilinear forms [R1]. The problem with sesquilinear forms is that one cannot

multiply them like matrices; if one tries to do that one gets in general a divergent expression. Langmann showed that there is a regularized multiplication of the forms which leads to the cocycle (6). In fact, one can work backwards from the algebra to construct the regularized product, [M2].

A computation of the vacuum expectation value (with respect to the free vacuum parametrized by $F = \epsilon$) of a product of operators $X_1 X_2 \dots X_n$ corresponding to charges (1) is now completely algebraic. Roughly speaking, any X_i can be split into an energy increasing operator X_+ and an energy lowering operator X_- . When acting on the vacuum on the right X_- gives zero whereas X_+ hitting the vacuum on the left in $\langle 0 | X_1 \dots X_n | 0 \rangle$ gives zero. Using repeatedly the commutation relations (6) one reduces the operator product to a product of the cocycles c_2 (which are functions of F) sandwiched between the vacua. As a final step one evaluates all cocycles at $F = \epsilon$, see [M2] for details.

In addition to the description of the current algebra as Fock bundle maps or as sesquilinear forms in a fixed Fock space there is a third alternative way for constructing a "representation". Let $\mathcal{F}_A$ be the family of (fermionic) Fock spaces parametrized by the space $\mathcal{A}$ of external vector potentials A . Since A is flat the Fock bundle $\mathcal{F}$ is trivial; choose a trivialization by choosing for every $A \in \mathcal{A}$ a unitary operator $h_A : H_0 \rightarrow H_A$, where H_A is the one-particle Hilbert space corresponding to the potential A . The operator h_A should satisfy the condition that $\epsilon_A h_A - h_A \epsilon_0$ is Hilbert-Schmidt; here $\epsilon_A = 2\pi_{+,A} - 1$, $\pi_{+,A}$ is the projection onto the positive energy subspace in the background A . If $S \in GL_1$ belongs to the unitary subgroup U_1 then also $h_A S$ satisfies the same condition. On the other hand, if both h_A and h satisfy the condition above, then $h = h_A S$ for some $S \in U_1$. It follows that the set of allowed h_A 's form a principal bundle P over $\mathcal{A}$ with structure group U_1 . Since the base space $\mathcal{A}$ is flat, the bundle has a smooth section h_A .

Denote $D_A^h = h_A^{-1} D_A h_A$. By the basic property of h_A the self-adjoint operator D_A^h is quantizable in the free fermionic Fock space $\mathcal{F}_0$, and for each λ not in the spectrum of D_A there is a unique vacuum ray $|A, \lambda\rangle$ characterized by

$$a^*(v)|A, \lambda\rangle = 0 = a(u)|A, \lambda\rangle$$

where v (resp. u) is an eigenvector of D_A^h belonging to an eigenvalue smaller (resp. bigger) than λ . Thus we have a trivial bundle of Fock spaces parametrized by the potentials. Note however that there is no continuous global section ψ such that $\psi(A)$ is a vacuum for D_A^h . This is because the eigenvalues of the Dirac operator, as a function of A , can cross any given vacuum level λ .

The action of a gauge transformation g is given as follows. Besides the usual gauge action $A \mapsto g \cdot A = g A g^{-1} + dg g^{-1}$ in the base space $\mathcal{A}$ we have an action in the fibers of the Fock bundle, which is obtained by quantizing the one-particle operator $T_A(g) = h_{g \cdot A}^{-1} T(g) h_A$. The quantization of the operator $T_A(g) \in GL_1$ is naturally defined only up to a phase (because of the nontrivial central extension $\widehat{GL}_1$). Again since $\mathcal{A}$ is flat, the phase for any given g can be chosen as a continuous function of A . However, as a function of g the choice cannot be continuous (topological obstruction from the nontriviality of $\widehat{GL}_1 \rightarrow GL_1$). If we have made a choice of the phase for three elements g_1, g_2 , and $g_1 g_2$, then we can write

$$\hat{T}_A(g_1 g_2) = \omega(g_1, g_2; A) \hat{T}_A(g_1) \hat{T}_A(g_2)$$

where the phase ω is also a function of the base point A .

Thus, although we still do not have a normal Hilbert space representation of the current algebra, we have a *cocycle valued in the representation of the smaller algebra $\widehat{GL}_1$* . Details of the construction will appear elsewhere.

In $1 + 1$ dimensions there is a famous "Bose-Fermi correspondence" based on representation theory of Kac-Moody algebras. There is a modification of the Bose-Fermi correspondence which works also in $3 + 1$ space-time dimensions. The first indication in this direction was the 'kink' construction of fermions by Finkelstein and Rubinstein, [FR]. For example, take $M = S^3$ and $G = SU(N)$. A kink is then a noncontractible map $f : S^3 \rightarrow G$. The homotopy classes of kinks are classified by an element of $\pi_3 G$, which is equal to $\mathbb{Z}$ when $N \geq 2$. When $N = 2$ the fourth homotopy group $\pi_4 G$ is $\mathbb{Z}_2$. In [FR] it was shown that in a configuration containing two nonoverlapping kinks a continuous rotation of one of the kinks is homotopic to a continuous interchange of the kink positions. This is a kind of spin-statistic theorem (later generalized by Sorkin, [So]). In particular, the case $G = SU(2)$ allows two types of field configurations. A contractible loop in $\mathcal{G} = \text{Map}(S^3, G)$ is interpreted as a boson whereas a noncontractible loop (generator of $\pi_4 SU(2)$) is a fermion.

The possibility of realization of fermions in terms of seemingly bosonic fields really occurs in the WZW model, as shown by Witten in [W]. The quantum mechanical WZW action functional obtains a factor -1 each time as the field $f : S^3 \rightarrow SU(n)$ is adiabatically rotated by the angle 2π . On the other hand, it was proved in [M3] that if $\mathcal{G}$ acts projectively in the quantum vector space (the nontrivial projective phases are related to chiral anomaly) then $D(f)D(g) = (-1)^{n(f)n(g)} D(g)D(f)$, where $n(f)$ is the winding number of f . I want to stress that this relation does not depend on the existence of a unitary (projective) representation D (that may not exist) in the ordinary sense. The group has nonunitary linear representations which have some similarities with the unitary highest weight representations in the one-dimensional case, [MR]. The formula is purely a result of the structural relations in an extension $\hat{\mathcal{G}}$.

A more recent and important result on the Bose-Fermi relation in $3 + 1$ dimensions is due to Ruijsenaars, [R2]. As we have seen, although the operators $D(g)$ representing gauge transformations do not exist as linear operators in the fermionic Fock space, they still exist as sesquilinear forms in a suitable dense domain. Furthermore, one can take limits of kinks $f_n : S^3 \rightarrow G$ such that $\lim D(f_n)$ exists as a bilinear in the Fock space, when $\lim f_n$ is a 'pointlike kink', the winding being concentrated to a single point $x \in S^3$. The limiting bilinear is the fermion field.

REFERENCES

- [FR] Finkelstein, D. and J. Rubinstein. J. Math. Phys. **9**, 1762 (1968). [FT] Fujii, K., and M. Tanaka. Commun. Math. Phys. **129**, 267 (1990). [L] Langmann, E. Proc. of the colloquium "Topological and Geometrical Methods in Field Theory", Turku, Finland, May 1991 (eds. by J. Mickelsson and O. Pekonen, World Scientific, Singapore, 1992). [Lu] Lundberg, L.-E. Commun. Math. Phys. **50**, 103 (1976). [M1] Mickelsson, J. Commun. Math. Phys. **127**, 285 (1990); [M2] ——— M.I.T. preprint CTP#2107, June 1992; [M3] ———: *Current Algebras and Groups*. Plenum Press, New York and London (1989). [MR] Mickelsson, J. and S. Rajeev. Commun. Math. Phys. **116**, 265 (1988). [R] Ruijsenaars, D. Commun. Math. Phys.

123, 617 (1989). [**R1**] Ruijsenaars, S.N.M. J. Math. Phys. **18**, 517 (1977); [**R2**]
——— Commun. Math. Phys. **124**, 553 (1989). [**So**] Sorkin, R. Commun. Math.
Phys. **115**, 421 (1988). [**W**] Witten, E. Nucl. Phys. **B223**, 433 (1983).